

# Problem B

## Average Substring Value

Time Limit: 1 second, Memory limit: 2G

Let  $s$  be a nonempty string consisting entirely of base-10 digits (0–9). If the length of  $s$  is  $n$ , number the digits  $1, 2, 3, \dots, n$  from left to right, and for  $1 \leq i \leq j \leq n$ , let  $s[i, j]$  denote the substring consisting of the digits from position  $i$  to position  $j$ , inclusive. (It follows that we are only considering *nonempty* substrings.) Assign a value to each substring that is simply equal to the largest digit in the substring. What is the average value of the substrings of  $s$ ?

Note that two different substrings may be identical (as strings), but for the purposes of this problem they are treated as distinct. For example, if  $s = 1010$ , then  $s[1, 2] = s[3, 4] = 10$  are distinct substrings (both with value 1).

### Input

The input is a single nonempty string,  $s$ , of base-10 digits. The length of  $s$  is at most 200 000.

### Output

Output a line containing the average value of the substrings of  $s$ . If the average is an integer, print the integer. If the average is a proper fraction, i.e., is equal to  $a/b$ , where  $a$  and  $b$  are positive integers and  $a < b$ , print this fraction in lowest terms, with a '/' symbol separating the numerator and denominator. If the average is greater than 1 and does not simplify to an integer, print the whole part followed by the proper fractional part, separated by a space, with the proper fractional part in lowest terms and formatted as described in the previous sentence.

#### Sample Input 1

123

#### Sample Output 1

2 1/3

#### Sample Input 2

4084

#### Sample Output 2

6

#### Sample Input 3

1010

#### Sample Output 3

4/5

#### Sample Input 4

00000

#### Sample Output 4

0

This page is intentionally left blank.